

## Epidemiology

Differential mortality risks associated with PM<sub>2.5</sub> components. A multi-country, multi-city study

Masselot, P; Sera, F; Schneider, R; Kan, H; Lavigne, E; Stafoggia, M; Tobias, A; Chen, H; Burnett, RT; Garland, Rebecca M

### **Abstract:**

**Background:** The association between fine particulate matter (PM<sub>2.5</sub>) and mortality widely differs between as well as within countries. Differences in PM<sub>2.5</sub> composition can play a role in modifying the effect estimates, but there is little evidence about which components have higher impacts on mortality. **Methods:** We applied a 2-stage analysis on data collected from 210 locations in 16 countries. In the first stage, we estimated location-specific relative risks (RR) for mortality associated with daily total PM<sub>2.5</sub> through time series regression analysis. We then pooled these estimates in a meta-regression model that included city-specific logratio-transformed proportions of seven PM<sub>2.5</sub> components as well as meta-predictors derived from city-specific socio-economic and environmental indicators. **Results:** We found associations between RR and several PM<sub>2.5</sub> components. Increasing the ammonium (NH<sub>4</sub><sup>+</sup>) proportion from 1% to 22%, while keeping a relative average proportion of other components, increased the RR from 1.0063 (95% confidence interval [95% CI] = 1.0030, 1.0097) to 1.0102 (95% CI = 1.0070, 1.0135). Conversely, an increase in nitrate (NO<sub>3</sub><sup>-</sup>) from 1% to 71% resulted in a reduced RR, from 1.0100 (95% CI = 1.0067, 1.0133) to 1.0037 (95% CI = 0.9998, 1.0077). Differences in composition explained a substantial part of the heterogeneity in PM<sub>2.5</sub> risk. **Conclusions:** These findings contribute to the identification of more hazardous emission sources. Further work is needed to understand the health impacts of PM<sub>2.5</sub> components and sources given the overlapping sources and correlations among many components.